

Poly- β -hydroxybutyrate Production by Root Nodule Isolates of Wild *Vicia faba*

Çiğdem Küçük* and Cenap Cevheri

Harran University, Faculty of Arts and Sciences, Department of Biology, Şanlıurfa, TURKEY

Received: 07.04.2016; Accepted: 25.05.2016; Published Online: 21.07.2016

ABSTRACT

Poly- β -hydroxybutyrate is a common bacterial polymer. It has been ecofriendly and best alternative biopolymer against environmental pollution. In this study, the poly- β -hydroxybutyrate (PHB) contents of *Rhizobium* isolates from isolated wild *Vicia faba* were studied in the media used by a spectrophotometric technique. The dry cell weights and PHB production of isolates were different. The production of PHB by 18 isolates ranged between 0.006-0.720 g/l with a productivity of 0.273-19.3 (%). B7 isolate produced more PHB than the other isolates.

Keywords: Poly- β -hydroxybutyrate, *Rhizobium*, isolate, wild *Vicia faba*

INTRODUCTION

The pastures are an important part of farming systems (Avcioğlu *et al.* 2009). Şanlıurfa's natural pastures are approximately 234.527 ha and *Vicia*, *Trifolium*, *Pisum sativum*, *Vicia faba* are the primary legumes in natural pasture in Şanlıurfa, Turkey (Cevheri and Polat 2009). The soil bacterium *Rhizobium* sp. fixes nitrogen in symbiotic association with the forage legumes (Hansen 1994). *Rhizobium* bacteria have the exceptional ability to form nodules on roots of leguminous plants (Hansen 1994, O'Hara *et al.* 2003). The use of *Vicia faba* as a forage legume in Turkey required the introduction of the corresponding symbionts. The green and dry herbs, straw, silage, residues of *Vicia faba* evaluated for animal feed. Also, because it has quite a green hitch it can be used as green manure crops. One of the biggest feature of *Vicia faba* is the crude protein content reaching 30% in some varieties. The protein digested ratio and biological value due to tannins, lectins, glycosides some inhibitory substances is low (Avcioğlu *et al.* 2009). The aim of the bean fodder production; as feed to make use of them. Also picks assessed before flowering or flowering process in order to meet the requirements of green feed ruminant animals (Cevheri and Polat 2009, Hansen 1994).

This species has wide adaptation and produces high quality forage under a grazing regimes (Avcioğlu *et al.* 2009). Many papers have reported enhancement of nodulation and growth of a wide variety of forage and grain legumes by *Rhizobium* bacteria (Hansen 1994, Seguin *et al.* 2001, Zahran 2001, Khanna and Srivastava 2005). Symbiotic nitrogen fixation is often limited as a result of environmental conditions such as nutrient deficiencies, soil pH, temperature, soil moisture, salinity (Hansen 1994, Zahran 2001).

Poly- β -hydroxybutyrate is the the most common microbial biodegradable plastics (Lee 1996). Poly- β -hydroxybutyrate (PHB) is synthesized under negative growth and environmental conditions by microorganisms such as bacteria, fungi and algae (Khanna and Srivastava 2005, Panigrahi and Badveli 2013). PHB are biodegradable polymers and they synthesized by numerous bacterial isolates such as root nodule bacteria (Anderson and Dawes 1990). Among these *Rhizobium* bacteria are *Rhizobium galega*, *R. heydsarum*, *R. leguminosarum*, *R. leguminosarum* biovar *trifolii*, *R. meliloti* (Tombilini and Nuti 1989, Tavernier *et al.* 1997).

These polymers are synthesized by *Rhizobium* bacteria when carbon source is abundant and growth is limited by the shortage of another nutrient (Zevenhuizen 1981). The purpose of this study was to isolate wild *Vicia faba* root nodule bacteria and determine PHB production by isolates.

* Corresponding author: ckucuk@harran.edu.tr

MATERIALS AND METHODS

Isolation

Rhizobium sp. isolates were isolated from root nodules of wild *Vicia faba* plants. For the isolation of *Rhizobium* sp. isolates from *Vicia faba*; uprooted roots having pink healthy nodules were selected, washed 5 times in distilled water and sterilized with 0.1% HgCl₂, crushed and grown in sterile yeast extract mannitol agar (Vincent 1970). Inoculated medium was incubated 28 °C for 5-6 days. The enriched sample was grown on congo red (0.0025 g/l of YEMA) containing Yeast Extract Mannitol Agar (YEMA) at 28 °C for 24 h. Isolates were purified and were tested for Gram reaction, colony structure, catalase, urease, growth at different pH levels, determination of NaCl tolerance, utilization of carbohydrates, hydrolysis of starch and gelatin (Jordan 1994). Acid and alkali production was determined in YEMA medium with bromothymol blue indicator (0.0025 %) (Jordan 1994).

Analytic method for PHB

Determination of the amount of chemically. The cultures were inoculated with a 2% (v/v) inoculum, grown for 48 h in Yeast Extract Mannitol Broth at 250 rpm for 28 °C. At the end of this period, the samples were centrifuged for 20 min at 6000 rpm. The pellets were suspended in 5 ml sterile distilled water and then homogenized. 2 ml of the culture suspension was added 2 ml of 2 M HCl, the contents were heated at boiling temperature for 2 h in a water bath. Then, the samples were centrifuged at 6000 rpm for 20 min. Five millilitre of chloroform were added to the resulting precipitate and the tubes were left overnight at 250 rpm at 28 °C. The contents were centrifuged at 6000 rpm for 20 min and extracted with 100 µl of chloroform, contents were dried at 60 °C. 10 ml 98 % sulphuric acid was added and heated at 60 °C for 1 h. The amount of PHB was measured at 235 nm in a UV spectrophotometer (Bonartseva and Myskina 1985, Bonartseva et al. 1994).

Also, cultures were grown at YEM broth, centrifuged at 6000 rpm for 30 min. The pellets were washed twice with sterile distilled water and dried for 24 h at 100 °C. The total bacterial dry weight was determined. All experiments were repeated three times. Their average values and Standard deviations (SD) are given.

RESULTS AND DISCUSSION

The bacteria isolated from the root nodules was identified as a *Rhizobium* sp. according to Bergey's Manual of Systematic Bacteriology (Jordan 1994) but the species was not ascertained. The bacteria were also identified following the methods given in the Manual of Microbiological Methods (Jordan 1994). The bacteria were rod shaped, aerobic, motile, gram negative and non-spore forming which showed positive results for catalase, acid production YEM agar, urease, citrate, nitrate reduction tests and could hydrolyze gelatin. The bacterial isolates showed negative results in citrate utilization tests, 4% NaCl tolerance tests. The bacterial isolates produced acid in sugar medium and serum zone in litmus milk medium. All of the isolates gave negative results for starch hydrolysis, indole, voges Proskauer and methyl red tests (Table 1).

Also, isolates have shown tolerance at 1 %, 2% and 3 % NaCl. Similarly, variation of *Rhizobium* isolates subjected to NaCl tolerance, growth at different pH levels and carbohydrates is reported (Panigrahi and Badveli 2013, Jordan 1994, Junior *et al.* 2011).

In this study, PHB production of the root nodule isolates were detected. The cell dry weight and yield of PHB produced by root nodule bacterial isolates are shown in Table 2. The dry cell weights of cultures were different (Table 2). The production of PHB by 18 isolates ranged between 0.006-0.720 g/l with a productivity of 0.27-19.3 (%).

Table 1. Some morphological and biochemical characteristics of isolates.

Characteristics		Number of positive isolates	Number of negative isolates
Gram reaction		18	-
Cell shape		Short rods	
Growth at	15 °C	6	12
	30 °C	18	-
	37 °C	10	8
Growth at pH	5	18	-
	7	18	-
	8	15	3
Acid production on YEMA		18	-
Alkali production on YEMA		-	18
NaCl tolerance	1 %	18	-
	2%	18	-
	3%	18	-
Utilization of carbohydrates	Mannose, mannitol, xylose, raffinose, glucose, galactose, fructose, arabinose	18	-
	dulcitol	-	18
Catalase, nitrate, urease, citrate,		18	-
Indole, methyl red, Voges prokauer		-	18
Hydrolysis	Starch	-	18
	Gelatin	18	-

Table 2. PHB production by root nodule isolates.

Isolates	Cell Dry Weight (g/l)	PHB (g/l)	Yield of PHB (%) [#]
B1	1.04 ± 0.04*	0.020 ± 0.003	1.92
B2	0.72 ± 0.110	0.016 ± 0.015	2.22
B3	0.80 ± 0.005	0.218 ± 0.012	0.27
B4	0.79 ± 0.160	0.082 ± 0.002	10.3
B5	0.27 ± 0.140	0.390 ± 0.004	1.44
B6	0.40 ± 0.03	0.445 ± 0.001	1.11
B7	1.18 ± 0.025	0.720 ± 0.021	6.10
B8	0.36 ± 0.035	0.006 ± 0.042	1.67
B9	0.65 ± 0.040	0.062 ± 0.006	9.53
B10	0.98 ± 0.050	0.013 ± 0.004	1.32
B11	0.72 ± 0.310	0.084 ± 0.005	11.7
B12	1.04 ± 0.025	0.092 ± 0.019	8.84
B13	0.57 ± 0.035	0.015 ± 0.015	2.63
B14	0.49 ± 0.050	0.095 ± 0.012	19.3**
B15	0.71 ± 0.620	0.017 ± 0.010	2.39
B16	0.65 ± 0.040	0.056 ± 0.018	8.61
B17	1.25 ± 0.052	0.050 ± 0.006	4
B18	1.17 ± 0.055	0.025 ± 0.001	2.13

*Values are the means ± Standard deviations of three measurements.

**The highest PHB production

[#]According to cell dry weight

B7 isolate produced more PHB than the other isolates and B8 produced less PHB than the other isolates did. In one of the the studies conducted by Hernandez *et al.* (2013), it was reported that *Rhizobium* isolates accumulated 0.001-0.009 g/l PHB during broth medium. The highest yield of PHB accumulation in comparison to dry weight was obtained in B14 isolate (19.3 %). Bergersen *et al.* (1991) reported that accumulate high levels of PHB during the active nitrogen fixing period of symbiosis of bean and soybean. Also, Rebah *et al.* (2009) reported that PHB yields (%) accumulated in cells according to dry weight were different; 7.27 % for *S.meliloti*; 5.23 % for *R.lupini*. Accumulated PHB in bacteroid can be mobilized for use during periods of low carbon availability (Rebah *et al.* 2009, Kretovich *et al.* 1977). Rebah *et al.* (2009), Lowing *et al.* (2005); Mercan *et al.* (2002) obtained that poly-β-hydroxybutyrates were important molecules in *Sinorhizobium meliloti*, *R.leguminosarum* bv. *phaseoli*, *Rhizobium leguminosarum* bv. *viciae*. PHB plays a role in cell protection during

stress periods such as drought, salt stresses, starvation and low nutrient availability (Tavernier *et al.* 1997, Tamdoğan and Sidal 2011).

In this paper, the PHB production by *Rhizobium*'s wild isolates were determined in the medium. Further work will be carried out to interactions between nitrogenase activity and PHB production.

REFERENCES

- Anderson AJ and Dawes EA (1990). Occurrence, metabolism, metabolic role, and industrial uses of bacterial polyhydroxyalkanoates. *Microbiol Rev.*, 54: 450-472.
- Avcıoğlu R, Hatipoğlu R and Karadağ Y (2009) Yem bitkileri T.C. Tarım ve Köyişleri Bak. Tarımsal Üretim ve Geliştirme Genel Müd. İzmir. s. 545 (in Turkish),
- Bergensen FJ, Peoples MB and Turner GL (1991). A role for poly- β -hydroxybutyrate in bacteroids of soybean root nodules. *Proceedings of the Royal Soc. B.*, 245: 59-64
- Bonartseva GA and Myskina VL (1985). Fluorescence intensity of strains of nodule Bacteria (*Rhizobium meliloti*, *R.phaseoli*) different in activity, grown in the presence of the lipophilic vital strain phosphine R. *Transl. of Microbiologiya.*, 54(4): 535-541.
- Bonartseva GA, Myskina VL and Zagreba ED (1994). Poly- β -hydroxybutyrate content in cells of various *Rhizobium* species during growth with different carbon and nitrogen sources. *Microbiol.*, 63(1): 45-48.
- Cevheri C and Polat T (2009). Şanlıurfa'da Yem Bitkileri Tarımının Dünü, Bugünü ve Yarını. Harran Üniversitesi Ziraat Fakültesi Dergisi (Journal of the Faculty of Agriculture) 13: 63-76. (in Turkish).
- Hansen AP (1994). Symbiotic N₂ Fixation of Crop Legumes: Achievement and Perspectives. Margraf Verlag, USA. pp. 31-36.
- Hernandez I, Mapoles MC, Rosales PR, Bonas R and Ramirez JF (2013). Selection of rhizobia isolates from nodules of the forage legume *Pueraria phaseoloides* (tropical kuzdu). *Cuban Journal of Agricultural Science*, 47: 311-318.
- Khanna S and Srivastava A (2005) Recent advances in microbial polyhydroxyalkanoates. *Process Biochem.* 40: 607-6019.
- Kretovich WL, Romanov VI, Yushkova LA, Shramko VI and Fedulova NG (1977). Nitrogen fixation and poly-B-hydroxybutyric acid content in bacteroids of *Rhizobium lupine* and *R.leguminosarum*. *Plant and Soil.*, 48:291-302.
- Lee SY (1996). Bacterial polyhydroxyalkanoates. *Biotechnol. Bioeng.*, 49: 1-14.
- Lowing EM, Leonard M, Marraoqui S, Wheeler TR, Findlay K, Downie JA and Poole PS (2005). Role of poly-hydroxybutyrate and glycogen as carbon storage compounds in pea and bean bacteroides. *Mol. Plant Microb. Interact.*, 18: 67-74.
- Mercan N, Aslım B, Yüksekdağ ZN and Beyatlı Y (2002). Production of poly- β -hydroxybutyrate by some *Rhizobium* bacteria. *Turk J Biol.*, 26:215-219.
- Jordan DC (1994). Family III. Rhizobiaceae. In: KRIEG, N.R. and Holt, J.G. eds. *Bergey's Manual of Systematic Bacteriology*, Willians and Wilkins, Baltimore, pp. 234-342.
- Junior PIF, de Oliveira PJ, Rumjanek NG and Xavier GR (2011). Poly- β -hydroxybutyrate and exopolysaccharide biosynthesis by bacterial isolates from pigeonpea (*Cajanus cajan* (L.) Mill sp.) root nodules. *Appl. Biochem. Biotechnol.*, 163: 473-484.
- O'Hara GW, Howieson JG, Graham PH (2003). Nitrogen fixation and agricultural practice. In: *Nitrogen Fixation in the Millennium* (ed. Leigh G.J.), pp. 391-410. Elsevier Press, London.
- Panigrahi S and Badveli U (2013). Screening, isolation and quantification of PHB-producing soil bacteria. *Int. J. of Eng. Sci. Innovation.*, 2(9): 1-6.
- Rebah FB, Prevost D, Tyagi RD and Belbahri L (2009). Poly- β -hydroxybutyrate production by fast growing Rhizobia cultivated in sludge and in industrial wastewater. *Appl. Biochem. Biotechnol.*, 158: 155-163.
- Seguin P, Graham PH, Sheaffer CC, Ehlke NJ and Russelle MP (2001). Genetic diversity of rhizobia nodulating *Trifolium ambiguum* in North America. *Canadian Journal of Microbiology*, 47: 81-85.
- Tamdoğan N, Sidal U (2011). Investigation of Poly- β -Hydroxybutyrate (PHB) production by *Bacillus subtilis* ATCC 6633 under different conditions. *Kafkas Univ Vet Fak Derg.*, 17: 173-176.
- Tavernier P, Portais JC, Nava Saucedo JE, Courtois J, Courtois B, Barbotin JN (1997). Exopolysaccharide and poly- β -hydroxybutyrate coproduction in two *Rhizobium meliloti* strains. *Appl Environ Microbiol.*, 63: 21-26.
- Tombilini R, Nuti MP (1989) Poly (β -hydroxyalkanoate) biosynthesis and accumulation by different *Rhizobium* species. *FEMS Microbiology Letters*, 60: 298-304.
- Vincent JM (1970). A manual for the practical study of root nodules bacteria. In *IBP handbook* (15th ed.). Oxford: Blackwell Scientific,
- Zahran HH (2001). Rhizobia from wild legumes: diversity, taxonomy, ecology, nitrogen fixation and biotechnology. *J Biotechnol.*, 91: 143-53.
- Zevenhuizen LPTM (1981). Cellular and extracellular polysaccharides in fast growing species of *Rhizobium*. *Antonie Van Leeuwenhoek*, 47: 481-497.